

INFORMATION DISCLOSURE STATEMENT PTO-1449	Atty. Docket No. 031729	Serial No. <sup>10/673,380</sup> <del>New Application</del>
	Applicant(s): NAKAGAWA, Takashi, et al.	
	Filing Date: September 30, 2003	Group Art Unit: <del>Unassigned</del> 1647

## U.S. PATENT DOCUMENTS

Examiner Initial	Document No.	Name	Date	Class	Subclass	Filing Date (If appropriate)
_____	AA					
_____	AB					

## FOREIGN PATENT DOCUMENTS

Document No.	Date	Country	Translation (Yes or No)
_____	AC		
_____	AD		

## OTHER DOCUMENTS

SLW		
_____	AE	Bertog, Marko, et al. "Basolateral proteinase-activated receptor (PAR-2) induces chloride secretion in M-1 mouse renal cortical collecting duct cells". <i>Journal of Physiology</i> (1999), 521-1, pp. 3 - 17.
_____	AF	Gui, Yu, et al. "Bidirectional regulation of renal hemodynamics by activation of PAR <sub>1</sub> and PAR <sub>2</sub> in isolated perfused rat kidney." <i>Am J Physical Renal Physiol</i> 285: F95-104, 2003.
_____	AH	Trottier, Greg, et al. "PAR-2 elicits afferent arteriolar vasodilation by NO-dependent and NO-independent actions." <i>Am J Physical Renal Physiol</i> 282: F891- 897, 2002.
_____	AI	Grandaliano, Giuseppe, et al. "Protease-Activated Receptor 2 Expression in IgA Nephropathy: A Potential Role in the Pathogenesis of Interstitial Fibrosis." <i>J Am Soc Nephrol</i> 14: 2072-2083, 2003.
_____	AJ	LeHir, Michel, et al. "IL-12-Dependent, IFN-γ-Independent Experimental Glomerulonephritis." <i>Kidney Blood Press Res</i> 2001; 24-27-32.
_____	AK	ITO, Mikio, et al. "Crescentic Type Nephritis Induced By Anti-Glomerular Basement Membrane (GBM) Serum in Rats." <i>Japan J. Pharmacol.</i> 33, 1145-1154 (1983).

SLW	AL	Kawabata, Atsufumi, et al. "Activation of Protease-Activated Receptor-22 (PAR-2) Triggers Mucin Secretion in the Rat Sublingual Gland." <i>Biochemical and Biophysical Research Communications</i> 270, 298-302 (2000).
	AM	Bahjat, Al-Ani, et al. "Detection of functional receptors for the proteinase-activated-receptor-2-activating polypeptide, SLIGRL-NH <sub>2</sub> , in rat vascular and gastric smooth muscle." <i>Can. J. Physiol. Pharmacol.</i> Vol. 73, 1203-1207 (1995).
	AN	Kong, Wuyi, et al. "Luminal trypsin may regulate enterocytes through proteinase-activated receptor 2." <i>Proc. Natl. Acad. Sci., USA</i> Vol. 94 pp. 8884-8889 (August 1997).
	AO	Molino, Maria, et al. "Endothelial Cell Thrombin Receptors and PAR-2." <i>The Journal of Biological Chemistry</i> Vol. 272, No. 17 pp. 11133-11141 (April 25, 1997).
	AP	Ferrell, William R., et al. "Essential role for proteinase-activated receptor 2 in arthritis." <i>The Journal of Clinical Investigation</i> Vol. 111, Number 1 (January 2003).
	AQ	Kawagoe, Junichi, et al. "Effect of Protease-Activated Receptor-2 Deficiency on Allergic Dermatitis in the Mouse Ear." <i>Japan J. Pharmacol.</i> 88, 77-84 (2002).
	AR	Lindner, Jonathan, et al. "Delayed Onset of Inflammation in Protease-Activated-Receptor-2-Deficient Mice." <i>The Journal of Immunology</i> pp. 6504-6510, Copyright 2000 by the American Association of Immunologists.
	AS	Kanke, Toru, et al. "Proteinase-activated Receptor-2 mediated Activation of Stress-activated Protein Kinases and Inhibitory $\kappa$ B Kinases in NCTC 2544 Keratinocytes." <i>The Journal of Biological Chemistry</i> , Vol. 276, No. 34, pp. 31657-31666 (Aug. 24, 2001).
	AT	Hou, L., et al. "Immunolocalization of protease-activated receptor-2 in skin: receptor activation stimulates interleukin-8 secretion by keratinocytes in vitro." <i>Immunology</i> 94, 356-362, Copyright 1998 Blackwell Science Ltd.
	AU	Compton, Steven J. et al. "The Role of Mast Cell Trypsin in Regulating Endothelial Cell Proliferation, Cytokine Release, and Adhesion Molecule Expression: Trypsin Induces Expression of mRNA for IL-1 $\beta$ and IL-8 and Stimulates the Selective Release of IL-8 from Human Umbilical Vein Endothelial Cells." <i>The Journal of Immunology</i> , pp. 1939-1946, Copyright 1998 by the American Association of Immunologists.
	AV	He, Shaoheng, et al. "Potent Induction of a Neutrophil and Eosinophil-Rich Infiltrate In Vivo by Human Mast Cell Trypsin: Selective Enhancement of Eosinophil Recruitment by Histamine." <i>The Journal of Immunology</i> , pp. 6216-6225; Copyright 1997 by the American Association of Immunologists.
	AW	Vergnolle, Nathalie "Proteinase-Activated Receptor-2-Activating Peptides Induce Leukocyte Rolling, Adhesion, and Extravasation in Vivo." <i>The Journal of Immunology</i> , pp. 5064-5069, Copyright 1999 by the American Association of Immunologists.

Received 9/30/2003

SLW	AX	Steinhoff, M., et al. "Agonists of proteinase-activated receptor 2 induce inflammation by a neurogenic mechanism." <i>Nature Medicine</i> Volume 6, Number 2 (February 2000)
	AY	He, Shaoheng and Walls, A. "Human mast cell tryptase: a stimulus of microvascular leakage and mast cell activation." <i>European Journal of Pharmacology</i> , 328 (1997) 89-97.
	AZ	Vergnolle, Nathalie, et al. "Characterization of the inflammatory response to proteinase-activated receptor-2 (PAR <sub>2</sub> )-activating peptides in the rat paw." <i>British Journal of Pharmacology</i> (1999) 127, 1083-1090.
	BA	Kawabata, Atsufumi, et al. "Increased vascular permeability by a specific agonist of protease-activated receptor-2 in rat hindpaw." <i>British Journal of Pharmacology</i> (1998) 125, 419-422.
	BB	Smith-Swintosky, et al. "Protease-Activated Receptor-2 (PAR-2) Is Present in the Rat Hippocampus and Is Associated with Neurodegeneration." <i>Journal of Neurochemistry</i> pp. 1890-1896 Copyright 1997 International Society for Neurochemistry.
	BC	Damiano, Bruce P., et al. "Increased Expression of Protease Activated Receptor-2 (PAR-2) in Balloon-Injured Rat Carotid Artery." <i>Thromb Haemost</i> 1999; 81:808-14.
	BD	Cicala, Carla, et al. "Protease-Activated Receptor-2 Involvement in Hypotension in Normal and Endotoxemic Rats in Vivo." <a href="http://www.circulationaha.org">www.circulationaha.org</a> . <i>Circulation</i> May 1999, Vol. 99: 2590-2597.
	BE	MacFarlane, Scott R., et al. "Proteinase-Activated Receptors." <i>Pharmacol Rev</i> 53:245-282, 2001.
	BF	Roy, Samir. S., et al. "Dual endothelium-dependent vascular activities of proteinase-activated receptor-2-activating peptides: evidence for receptor heterogeneity." <i>British Journal of Pharmacology</i> (1998) 123, 1434-1440.
	BG	Matsui-Nakanishi, Mayumi, et al. "PAR3 is a cofactor for PAR4 activation by thrombin." <i>Nature</i> vol 404, pp. 609-613 April 6, 2000.
	BH	Nystedt, Sverker, et al. "Molecular cloning of a potential proteinase activated receptor." <i>Proc. Natl. Acad. Sci. USA</i> Vol. 91, pp. 9208-9212, September 1994.

Examiner

/Sandra Wegert/

Date Considered

01/30/2007

<sup>1</sup>Applicant's unique citation designation number (optional). <sup>2</sup>See Kind Codes of USPTO Patent Documents at [www.uspto.gov](http://www.uspto.gov), MPEP 901.04 or in the comment box of this document. <sup>3</sup>Enter Office that issued the document, by the two-letter code (WIPO Standard ST. 3). <sup>4</sup>For Japanese patent documents, the indication of the year of the reign of the Emperor must precede the serial number of the patent document. <sup>5</sup>Kind of document by the appropriate symbols as indicated on the document under WIPO Standard ST. 16 if possible. <sup>6</sup>Applicant is to indicate here if English language Translation is attached.